


PATENT SPECIFICATION

(11) 1 464 005

1 464 005

- (21) Application No. 17492/74 (22) Filed 22 April 1974
 (23) Complete Specification filed 3 April 1975
 (44) Complete Specification published 9 Feb. 1977
 (51) INT CL^a A01N 9/20
 (52) Index at acceptance
 ASE 1A3G 1A5B2 1C14 1C15A3 1C15A5 1C15B3 1C15C1
 1C15E 1C15F3 1C5J 1C5K 1C5P
 (72) Inventors LIONEL VICTOR ALLAN and
 ANTHONY JOHN HINTON



SCIENCE REFERENCE LIBRARY

(54) METHOD FOR THE DESTRUCTION OF AQUATIC MOLLUSCS

(71) We, IMPERIAL CHEMICAL INDUSTRIES LIMITED, Imperial Chemical House, Millbank, London SW1P 3JF, a British Company, do hereby declare the invention, for which we pray that a patent may be granted to us, and the method by which it is to be performed, to be particularly described in and by the following statement:—

This invention relates to a method for the destruction of aquatic molluscs, particularly those molluscs which are responsible for the transmission of Bilharziasis.

Bilharziasis is a disease which is endemic in parts of Africa, the Middle East and the Far East. It is a debilitating disease and results in prolonged ill-health due to damage inflicted on various internal organs, particularly the liver. Bilharziasis is water-borne, and an essential element in its transmission is the function of certain aquatic snails as hosts for the organisms which are responsible for the disease. Human infection occurs under insanitary conditions; healthy people become infected by, for example, bathing in infected water, and water can become infected as a result of insanitary practices by infected individuals.

The disease can be attacked in three ways. Firstly, infected individuals can be treated with various drugs to eliminate the infection. Secondly, sanitary conditions can be improved in order to avoid transmission of the disease. Thirdly, the life-cycle of the organisms responsible for the disease can be interrupted by destruction of the snail hosts. It is with this last approach that the present invention is concerned.

Compounds which have previously been used as aquatic molluscicides include copper salts, especially copper sulphate, salts of pentachlorophenol, salicylic acid derivatives such as 5 - chlorosalicylic acid and dichloro-nitrosalicylic acid, N - triylmorpholine and zinc dialkyldithiocarbamates.

At a concentration which is lethal to aquatic snails, copper sulphate is not harmful to man or animals. However, especially in adverse

water conditions, i.e. turbid water containing much suspended matter, a concentration as high as 30 parts per million may be necessary, because some of the copper sulphate is rendered ineffective by precipitation under certain pH conditions and/or is removed by organic material. The necessity for clearing vegetation from the water to be treated thus raises the cost of the treatment. Copper sulphate is also corrosive towards metals, i.e. the pipes, tanks, pumps, etc., which are used to meter it into the water being treated. Because of the high concentration required, large quantities of the chemical have to be transported, again raising the cost of this method of treatment.

Sodium pentachlorophenate is also widely used, and at molluscicidal strength is very toxic to snail eggs as well as to the snails themselves, although not to man or animals. It is not seriously affected by the presence of vegetation, and the concentration necessary to kill the snails in flowing water conditions is about 10 p.p.m. for 8 hours. It is cheap to use, but solid sodium pentachlorophenate is a powerful irritant, and both solid and concentrated aqueous solutions thereof are highly toxic, so that adequate handling precautions must be taken.

Copper pentachlorophenate, prepared *in situ* by mixing copper sulphate and sodium pentachlorophenate in the water to be treated, is essentially insoluble in water and is consequently much more persistent than either copper sulphate or sodium pentachlorophenate individually. It is an effective molluscicide, but the unpleasant effects of handling toxic chemicals have still to be considered and precautions taken.

5 - chlorosalicylic acid is lethal to aquatic snails at about 1 p.p.m. and also kills snails' eggs, but is not toxic to mammals at this concentration. It is sparingly soluble in water. Undesirable features are that it may scorch vegetation and destroy algae. The compound is destroyed by the action of ultra-violet light.

All the molluscicides in current use have certain disadvantages and at their effective

50

55

60

65

70

75

80

85

90


95

concentrations as molluscicides may also have an adverse effect on desirable aquatic life such as vegetation and fish.

It has now been found that certain polymeric biguanides are effective aquatic molluscicides.


According to the present invention there is provided a method for the destruction of

aquatic molluscs and/or their eggs in water infested therewith which comprises adding to the aquatic habitat of the molluscs and/or their eggs to provide a concentration from 0.1 to 100 parts per million, a linear polymeric biguanide or a salt thereof which in its free base form has a recurring polymer unit represented by the formula



wherein X and Y represent bridging groups in which together the total number of carbon atoms directly interposed between the pairs of nitrogen atoms linked by X and Y is more than 9 and less than 17.


The bridging groups X and Y may consist of polymethylene chains, optionally interrupted by hetero atoms, for example, oxygen, sulphur or nitrogen. X and Y may also incorporate cyclic nuclei which may be saturated or unsaturated, in which case the number of carbon atoms directly interposed between the pairs of nitrogen atoms linked by X and Y is taken as including that segment of the cyclic group, or groups, which is the shortest. Thus, the number of carbon atoms directly interposed between the nitrogen atoms in the group



is 4 and not 8.

The preferred polymeric biguanide for use in the present invention is poly(hexamethylene biguanide), in which X and Y both represent the $-(CH_2)_6-$ group. The compound is preferably employed in the form of its hydrochloride salt, which is conveniently used as a 20% w/w aqueous solution (i.e. 100 parts by weight of the solution contain 20 parts by weight of the active agent). Other salts of polymeric biguanides using organic and inorganic acids may be used, for example, the sulphate, acetate, gluconate or behenate.

Polymeric biguanides may be prepared by the reaction of a bisdicyandiamide having the formula




with a diamine $H_2N-Y-NH_2$, wherein X and Y have the meanings defined above; or

by reaction between a diamine salt of dicyanamide having the formula



with a diamine $H_2N-Y-NH_2$, wherein X and Y have the meanings defined above. These methods of preparation are described in U.K. Specifications Nos. 702268 and 1152243 respectively, and any of the polymeric biguanides described therein may be used according to the present invention.


The polymer chains are terminated either by an amino group or salt thereof or by an




group, and the terminating groups may be the same or different on each polymer chain.

A small proportion of the primary amine $R-NH_2$, where R represents an alkyl group containing from 1 to 18 carbon atoms, may be included with the diamine $H_2N-Y-NH_2$ in the preparation of polymeric biguanides as described above. The primary monoamine acts as a chain-terminating agent and consequently one or both ends of the polymeric biguanide polymer chains may be terminated by an $-NHR$ group. The use of these chain-stopped polymeric biguanides is also within the scope of the present invention.

It is to be understood that by either of the above-described methods the polymeric biguanides are obtained as mixtures of polymers in which the polymer chains are of different lengths, the number of individual biguanide units, i.e.




and



together being from 3 to about 80.

the exposure period of the snails: the biocide solution was extended to 4 days, renewing the test solution each day. Dosages of 2, 1, 0.5, 0.25, 0.1 and 0.05 p.p.m. of poly(hexamethylene biguanide) hydrochloride were used. The LC_{50} for adult snails after 4 days was 0.25–0.5 p.p.m.



For 1—2 day old egg masses (the eggs hatch in 6—7 days under laboratory conditions) 20 p.p.m. of poly(hexamethylene biguanide) hydrochloride were required to kill embryos using a 24 hour exposure. However, with a 4 day exposure, the control groups and those exposed to 0.1 and 0.05 p.p.m. of the biocide hatched when they were 7 days old. With higher concentrations, the following results were obtained:

2.0 p.p.m.—all embryos died in the egg mass within 11 days after the treatment period.


1.0 p.p.m.—all embryos died in the egg mass within 21 days after the treatment period.

0.5 p.p.m.—some hatchlings occurred during the 21 day recovery period but the hatchlings did not appear to feed and eventually died.

0.25 p.p.m.—though hatching was delayed, about 75% of the embryos had hatched after 21 days and these fed and began growing.

It is clear from the above results that as little as 0.5 p.p.m. of poly(hexamethylene biguanide) hydrochloride is effective in killing the embryos of *Biomphalaria glabrata* after a 4 day exposure period.

1. A method for the destruction of aquatic molluscs and/or their eggs in water infested therewith which comprises adding to the aquatic habitat of the molluscs and/or their eggs to provide a concentration from 0.1 to 100 parts per million by weight, a linear polymeric biguanide or a salt thereof which in its free base form has a recurring polymer unit represented by the formula:



the polymeric biguanide is poly(hexamethylene biguanide) hydrochloride.

3. A method as claimed in claim 1 or claim 2 wherein the polymeric biguanide is employed at a concentration of approximately 1 part per million.

4. A method for the destruction of aquatic

molluscs and/or their eggs in water infested
therewith, substantially as hereinbefore de-
scribed with reference to the foregoing
Example.

D. VINCENT,
Agent for the Applicant.

Printed for Her Majesty's Stationery Office, by the Courier Press, Leamington Spa, 1977
Published by The Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from
which copies may be obtained.